

Freshwater Calanoida of British Columbia (June 03, 2014)

compiled by Ian Gardiner

This list is based primarily on Sandercock & Scudder (1996), and also on the additional references below.

Taxonomy is in accordance with the World of Copepods Taxonomic Database (Calanoida) available at

<http://www.marinespecies.org/copepoda/aphia.php?p=taxdetails&id=1100>

A number of other species (including *Arctodiaptomus arapahoensis*, *Hesperodiaptomus shoshone*, *Leptodiaptomus cuauhtemoci*, *Leptodiaptomus judayi*, *Osprhanticum labronectum*) are represented as occurring in British Columbia in several lists and keys. It is quite likely that some of these do indeed occur in British Columbia, but I have been unable to substantiate this by reference to unambiguous museum collections or unambiguous collections reported in peer-reviewed journals (ambiguity typically occurring in older collections taken at a time when some modern-day species were conflated with each other).

Family AETIDEIDAE Giesbrecht, 1892

Genus *Senecella* Juday, 1923
Senecella calanoides Juday, 1923

Family DIAPATOMIDAE Baird, 1850

Genus *Acanthodiaptomus* Kiefer, 1932
Acanthodiaptomus denticornis (Wierzejski, 1887)

Genus *Aglaodiaptomus* Light, 1938
Aglaodiaptomus forbesi Light, 1938
Aglaodiaptomus leptopus (Forbes S.A., 1882)

Genus *Hesperodiaptomus* Light, 1938
Hesperodiaptomus arcticus (Marsh, 1920)
Hesperodiaptomus caducus Light, 1938
Hesperodiaptomus eiseni (Lilljeborg, 1889)
Hesperodiaptomus franciscanus (Lilljeborg, 1889)
Hesperodiaptomus hirsutus Wilson M.S., 1953
Hesperodiaptomus kenai Wilson M.S., 1953
Hesperodiaptomus kiseri (Kincaid, 1953)
Hesperodiaptomus nevadensis Light, 1938
Hesperodiaptomus novemdecimus Wilson M.S., 1953

Genus *Leptodiaptomus* Light, 1938
Leptodiaptomus angustilobus (Sars G.O., 1915)
Leptodiaptomus ashlandi (Marsh, 1893)
Leptodiaptomus connexus Light, 1938
Leptodiaptomus insularis (Kincaid, 1956)
Leptodiaptomus minutus (Lilljeborg, 1889)

Leptodiaptomus novamexicanus (Herrick, 1895)
Leptodiaptomus nudus (Lilljeborg, 1889)
Leptodiaptomus sicilis (Forbes S.A., 1882)
Leptodiaptomus siculooides (Lilljeborg, 1889)
Leptodiaptomus signicauda (Lilljeborg, 1889)
Leptodiaptomus tyrelli (Poppe, 1888)

Genus *Onychodiaptomus* Light, 1939
Onychodiaptomus hesperus Wilson M.S. & Light, 1951
Onychodiaptomus sanguineus (Forbes S.A., 1876)

Genus *Skistodiaptomus* Light, 1939
Skistodiaptomus oregonensis (Lilljeborg, 1889)
Skistodiaptomus pallidus (Herrick, 1879)

Family TEMORIDAE Giesbrecht, 1893

Genus *Epischura* Forbes S.A., 1882
Epischura nevadensis Lilljeborg, 1889

Genus *Eurytemora* Giesbrecht, 1893
Eurytemora affinis (Poppe, 1880)

Genus *Heterocope* Sars G.O., 1863
Heterocope septentrionalis Juday & Muttkowski, 1915

References

Chengalath, R. & C. Shih 1994. Littoral freshwater copepods of northwestern North America: Northern British Columbia. *Verhandlungen Internationale Vereinigung für Theoretische und Angewandte Limnologie* 25: 2421-2431.

Sandercock, G.A. & G.G.E. Scudder 1996. *Key to the Species of Freshwater Calanoid Copepods of British Columbia*. Resources Inventory Committee, Province of British Columbia: vi + 51.

The above document (with older taxonomy) is available as part of a 1994 unpublished work on BC calanoid copepods by the same authors at

http://www.for.gov.bc.ca/hts/risc/o_docs/aquatic/calanoid/assets/final.pdf

Shih, C. & R. Chengalath 1994. Freshwater littoral copepods of British Columbia and Alberta (Canada). *Hydrobiologia* 292/293: 379-388.